

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/44286 A2

(51) International Patent Classification⁷: C07K 14/00

(21) International Application Number: PCT/US00/34194

(22) International Filing Date:
15 December 2000 (15.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/171,042 16 December 1999 (16.12.1999) US
60/234,572 22 September 2000 (22.09.2000) US

(71) Applicant (for all designated States except US): WHITE-
HEAD INSTITUTE FOR BIOMEDICAL RESEARCH
[US/US]; Nine Cambridge Center, Cambridge, MA 02142
(US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ROOT, Michael, J.
[US/US]; 10 Emerson Place, Apartment 12C, Boston, MA

02114 (US). KAY, Michael, S. [US/US]; 884 Main Street,
Apartment 3F, Cambridge, MA 02139 (US). CHAN,
David, C. [US/US]; 205 Kent Street #38, Brookline, MA
02446 (US). KIM, Peter, S. [US/US]; 48 Baskin Road,
Lexington, MA 02420 (US).

(74) Agents: COLLINS, Anne, J. et al.; Hamilton, Brook,
Smith & Reynolds, P.C., Two Militia Drive, Lexington, MA
02421 (US).

(81) Designated States (*national*): CA, JP, US.


(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: FIVE-HELIX PROTEIN



(57) Abstract: Five-Helix protein, which comprises the three N-helices and at least two, but not three, of the C-helices of the trimer-of-hairpin structure of HIV gp41, separated by linkers, such as amino acid residue linkers, is disclosed. Six-Helix protein, which includes the three N-helices and the three C-helices of the trimer-of-hairpin structure of HIV gp41, separated by linkers, is also disclosed.

WO 01/44286 A2

AL2

FIVE-HELIX PROTEIN

RELATED APPLICATION(S)

This application claims the benefit of U.S. Provisional application 60/171,042, entitled "Five-Helix Protein," by Michael J. Root, Michael S. Kay, David C. Chan and Peter S. Kim (filed December 16, 1999) and U.S. Provisional application 60/234,572, entitled "Protein Design of an HIV Entry Inhibitor," by Michael J. Root, Michael S. Kay, David C. Chan and Peter S. Kim (filed September 22, 2000). The entire teachings of both of the referenced provisional applications are incorporated herein by reference.

10 GOVERNMENT SUPPORT

The invention was supported, in whole or in part, by Grant Number PO1 GM 56552 from National Institutes of Health. The Government has certain rights in the invention.

BACKGROUND OF THE INVENTION

15 HIV is the virus that is responsible for the worldwide AIDS epidemic. The initial stages of HIV infection involve the fusion of the viral membrane with the target cell membrane, a process that injects the viral contents into the cellular cytoplasm. On the viral side, the molecular complex responsible for the fusion activity contains the surface protein gp120 and the transmembrane protein gp41. It is
20 currently believed that gp120 interacts with the proteins CD4 and coreceptors on the target cell, resulting in a conformational change that causes gp41 to insert its amino terminus (fusion peptide region) into the target cell membrane. This structural rearrangement promotes the fusion of virus and cellular membranes through a poorly understood mechanism.

SUMMARY OF THE INVENTION

The present invention relates to a novel protein, referred to as Five (5)-Helix or Five-Helix protein, that, under the conditions described herein, folds into a stable structure, binds a peptide (referred to as C34) that corresponds to the C-peptide region of HIV gp41 protein or a portion of the region and inhibits HIV infection of mammalian cells, such as human cells. Five-Helix is made up of the three N-helices and at least two, but not three, of the three C-helices of the trimer of hairpin structure of HIV gp41, separated by linkers, such as amino acid residue linkers. That is, Five-Helix includes the three N-helices and at least two of the three C-helices of HIV gp41. It can also include a portion of the third C-helix, but does not include the entire third C-helix. In each case, the helices are separated by linkers, preferably amino acid residue linkers, between the preceding and following helices. In one embodiment, Five-Helix can be represented as: N-linker-C-linker-N-linker-C-linker-N, wherein N represents an N-helix and C represents a C-helix or C-helix portion.

As used herein, the term Five-Helix or Five-Helix protein encompasses all such embodiments (those including three N-helices and two or more, but less than three complete C-helices, separated by appropriate linkers). The amino acid composition of Five-Helix can vary greatly, provided that Five-Helix presents a surface that is structurally complementary to the C-peptide region of HIV gp41 protein and, preferably, binds C34 or the C-peptide region of gp41, as peptides or part of gp41 as a whole. That is, the remaining (interacting) surface of Five-Helix (the C-peptide binding site, all or a portion of which is not occupied by a C-peptide) must be presented in such a manner (conformation) that it is available to bind the C-peptide region of HIV gp41. In the case of vaccine and therapeutic applications of Five-Helix, Five-Helix must bind (be capable of binding) C34 or the C-peptide region of HIV gp41. In the cases in which Five-Helix is used as a drug-screening tool or an antibody-screening tool, Five-Helix need not bind (need not be capable of binding) C34 or the C peptide region of HIV gp41.

In one embodiment, Five-Helix has the amino acid sequence of SEQ ID NO.: 1. In other embodiments, Five-Helix presents a surface that is structurally complementary to the C-peptide region, preferably binds C34 or the C-

peptide region and has an amino acid sequence that differs from that of SEQ ID NO.: 1 by addition, deletion, substitution or alteration of at least one amino acid residue. The order of the N-helices and C-helices of Five-Helix can also vary, provided that the conformation is such that the exposed protein presents a surface
5 structurally complementary to the C-peptide region of HIV gp41. The linkers can be of any length or composition, provided that the Five-Helix protein conformation, described above, is retained. Five-Helix can be an L-amino acid protein, a D-amino acid protein or a combination of L-amino acid residues and D-amino acid residues; these residues can be modified residues.

10 The present invention further relates to DNA encoding Five-Helix; methods of producing Five-Helix; methods in which Five-Helix is used, such as in methods of inhibiting entry of HIV into mammalian cells, including human cells, and methods of eliciting an immune response in an individual, such as a human; methods in which DNA encoding Five-Helix is used, such as in gene therapy methods;
15 genetically engineered cells, such as bacteria, human and other mammalian and other eukaryotic cells, which contain and express Five-Helix protein-encoding DNA and methods of using such cells (e.g., for gene therapy or Five-Helix production); compositions, such as pharmaceutical compositions, which include Five-Helix; Five-Helix complex comprising Five-Helix and a component that binds HIV
20 envelope protein (e.g., gp120); compositions, such as pharmaceutical compositions, which include Five-Helix complex; antibodies, particularly neutralizing antibodies which bind Five-Helix and methods in which such antibodies are used, such as methods of reducing HIV infection; and methods of identifying molecules or compounds that inhibit HIV infection of cells and/or bind the Five-Helix protein.

25 Five-Helix is useful as an anti-HIV therapeutic agent, a prophylactic agent or drug to prevent HIV infection, a reagent for identifying (screening for) or designing other anti-HIV therapeutics or prophylactics, and an immunogen to elicit antibodies that prevent or reduce HIV infection. In a specific embodiment, the invention relates to a method of identifying a compound or molecule that binds Five-Helix and
30 inhibits HIV infection of mammalian cells, wherein the compound or molecule to be assessed is referred to as a candidate inhibitor, comprising combining a candidate

-4-

inhibitor and Five-Helix, under conditions appropriate for binding of an inhibitor and Five-Helix to occur and determining if binding occurs, wherein if binding occurs, the candidate inhibitor is a compound or molecule that binds Five-Helix. The method optionally further comprises determining if the compound or molecule
5 that binds Five-Helix inhibits HIV infection of mammalian (e.g., human) cells, such as in a cell-based assay. Such a compound or molecule will inhibit (totally or partially) HIV infection of cells (e.g., by preventing or interfering with formation of the trimer-of-hairpins).

In another embodiment, the invention relates to a method of eliciting an
10 immune response to HIV in an individual, comprising introducing, by an appropriate route, a composition comprising Five-Helix and a physiologically acceptable carrier, in a dose sufficient to elicit the immune response in the individual. Vaccines comprising Five-Helix (or a variant or portion thereof) in a physiologically acceptable carrier are the subject of this invention.

15 Also the subject of the present invention is Six (6)-Helix protein, which comprises three N-helices and three C-helices of HIV gp41, joined by linkers, such as amino acid residue linkers. In one embodiment, Six-Helix protein comprises the amino acid sequence of SEQ ID NO.: 2. In other embodiments, the amino acid sequence of Six-Helix differs from that of SEQ ID NO.: 2 by addition, deletion,
20 substitution or alteration of at least one amino acid residue. Six-Helix protein is useful not only for producing Five-Helix, but also as a negative control in screening for drugs that inhibit membrane fusion.

BRIEF DESCRIPTION OF THE DRAWINGS

The file of this patent contains at least one drawing in color. Copies of this
25 patent with color drawing(s) will be provided by the Patent and Trademark Office upon request and payment of the necessary fee.

Figures 1A and 1B illustrate targeting HIV-1 membrane fusion. Figure 1A is a schematic of HIV-1 membrane fusion depicting events that promote formation of the gp41 trimer-of-hairpins. The N-terminal fusion peptide of gp41 (red),

-5-

inaccessible in the native state, inserts into target cell membranes following gp120 interaction with CD4 and coreceptors (not shown). Formation of the prehairpin intermediate exposes the N-terminal coiled coil (gray), the target of C-peptide inhibition. This transient structure collapses into the trimer-of-hairpins state that
5 brings the membranes into close apposition for fusion. Figure 1B shows the design of the 5-Helix construct. The ribbon diagrams (top) depict the core structure of the trimer-of-hairpins (left and D.C. Chan et al., Cell 89, 263-273 (1997)) and a model of 5-Helix (right). The inner gray helices represent N36 peptides, and the outer blue helices represent C34 peptides. One C-peptide has been removed in the model of 5-
10 Helix and orange lines have been drawn to represent connectivity between the helices. In the design of 5-Helix, the N40 and C38 sequences (given in single-letter amino acid code) are alternately linked by short Gly/Ser peptide sequences (gray bars in schematic at bottom (See Example 1).

Figures 2A – 2D show properties of 5-Helix. Figure 2A is the circular
15 dichroism (CD) spectrum of 5-Helix (10 μ M) at 25 $^{\circ}$ C. The spectrum indicates that the 5-Helix protein adopts >95% of the helical content expected from the design. Figure 2B is a graphic representation of thermal denaturation of 5-Helix monitored by ellipticity at 222 nm in TBS (filled squares) and in 3.7 M guanidine
(Gu)HCl/TBS (open squares). The denaturation observed in the GuHCl solution is
20 >90% reversible. Figure 2C shows results of nickel (Ni)-NTA precipitation of 5-Helix with a His-tagged C-peptide. Untagged 5-Helix and His-tagged C-peptide (denoted C37-H6) were mixed before Ni-NTA agarose was added in order to precipitate complexes containing C37-H6 (lanes 1 and 5 and Example 4). Addition of excess untagged C-peptide (C34) shifts the 5-Helix molecules from the bound to
25 the unbound fraction (lanes 2 and 6). Figure 2D is the CD spectra of 5-Helix and C37-H6 before (filled squares) and after (open circles) mixing in a mixing cuvette. The increase in ellipticity at 222 nm upon mixing indicates an interaction between the two species that increases the total helical content (corresponding to an additional 28 helical residues per associated C-peptide).

30 Figures 3A-3C show results of assessment of 5-Helix inhibition of HIV-1 envelope-mediated membrane fusion, as described in Example 2. Figure 3A shows

results of assessment of titration of viral infectivity by 5-Helix (filled squares), 6-Helix (open triangles), and 5-Helix(D4) (open circles), as described in Example 3 and 5. The data represent the mean \pm SEM of two or more separate experiments. Figure 3B is a graphic representation of antagonistic inhibitory activities of 5-Helix and C34. The number of syncytia were measured in a cell-cell fusion assay performed in the absence or presence of 5-Helix, C34, or mixtures of 5-Helix and C34 at the indicated concentrations. The IC_{50} values for 5-Helix and C34 in this assay are 13 ± 3 nM and 0.55 ± 0.03 nM, respectively (D. C. Chan et al., Proc. Natl. Acad. Sci. USA 95, 15613-15617 (1998)). Data represent the mean and range of mean of duplicate measurements, except for the control (mean \pm SEM of five measurements). Figure 3C shows results of assessment of 5-Helix inhibition of pseudotyped virus containing different HIV-1 envelope glycoproteins. The reported IC_{50} values represent the mean \pm SEM of three independent experiments.

Figure 4 is a helical wheel diagram depicting the interaction of 5-Helix with the C-peptide region of gp41. The a through g positions in each helix represent sequential positions in the 4,3-hydrophobic heptad repeat in each sequence. The a and d positions in the gp41 C-peptide region interact with the exposed e and g positions on the N40 coiled coil of 5-Helix. Residues are boxed according to their degree of conservation as determined from the alignment of 247 sequences from HIV-1, HIV-2, and SIV isolates (HIV-1 sequence database, August 2000, Los Alamos National Laboratory): black rectangle, >90% identical; grey rectangle, >90% conservative substitution; dotted rectangle, 70-90% conserved; no box, <70% conserved. In generating Figure 4, substitutions within the following groups of amino acid residues were considered to be conservative: [Asp, Glu], [Lys, Arg], [Asn, Gln], [Phe, Tyr], [Ser, Thr] and [Val, Ile, Leu, Met]. Note the high degree of conservation in the a and d positions of the C-peptide region of gp41, a property markedly lacking in other positions (particularly c and g) of the C-peptide region not directly involved in binding 5-Helix.

Figure 5 is the structural arrangement of HIV gp41. Helical regions (heptad repeats) are shown in grey, and the relative position of N- (N36) and C- (C34,

-7-

DP178) peptides are indicated. In the ribbon diagram of the helical region, the N-peptides are in light grey, while the C-peptide are in dark grey.

Figure 6 is the sequence of 6-Helix and 5-Helix. The predicted helical segments are designated by the stacked sequence.

5 Figure 7 is a ribbon diagram of one of the possible α -helical arrangements of 5-Helix. The N-helical trimer is light grey, the C-helical regions are in dark grey and the extended loop regions are in black (based on the structure of D.C. Chan, et al. (Cell 89, 263-273 (1997))).

10 Figure 8 shows images of cell-cell fusion assay titration experiments. The syncytia (representing fused cells) are blue in the image while debris is brown.

Figure 9 shows images of cell-cell fusion assay competition experiments. The amount of syncytia are recorded for cultures incubated in 200 nM 6-Helix or 5-Helix with increasing amounts of C34 peptide.

15 Figure 10 is a schematic of the design of the Five-Helix constructs. The schematic diagram depicts the linkage pattern of the basic 5-Helix construct. Three different C-termini were added. In 6-Helix, a His-tagged C-peptide is attached to 5-Helix in order to mimic the complete six-helix bundle of the trimer-of hairpins. The N40 and C38 sequences (alternately joined using short Gly/Ser linkers) are derived from the N- and C-peptide regions of HIV HXB2 gp41. (The red- and blue-boxed
20 residues depict the sequences of the N36 (SEQ ID NO.: 11) and C34 (SEQ ID NO.: 12) peptides, respectively.)

Figures 11A-11C show amino acid sequences of peptides (SEQ ID NOS.: 1-10) of the present invention.

25 DETAILED DESCRIPTION OF THE INVENTION

The conformation of a major part of the ectodomain of the gp41 molecule consists of a trimer-of-hairpins structure. The core "trimer-of-hairpins" is comprised of a central three-stranded N-helix coiled coil surrounded by three outer C-helices, forming a bundle with a total of six helices. The trimer-of-hairpins is a common
30 structural element involved in the fusion of many enveloped viruses, suggesting a critical role for this motif in promoting membrane fusion. In HIV gp41, the core of

the trimer-of-hairpins is a bundle of six α -helices (formed by the C-terminal regions of three gp41 ectodomains) packed in an antiparallel manner against a central, three-stranded coiled coil (formed by the N-terminal regions of the gp41 molecules) (M. Lu et al., J. Mol. Biol. 290, 1031-1044 (1995); D.C. Chan et al., Cell 89, 263-273 (1997); W. Weissenhorn et al., Nature 387, 426-430 (1997)); K. Tan et al., Proc. Natl. Acad. Sci. USA 94, 12303-12308 (1997). Because the fusion peptide region, which inserts into the cellular membrane, is located at the extreme N-terminus of gp41, and the C-terminal region is adjacent to the transmembrane helix anchored in the viral membrane, the trimer-of-hairpins motif serves to bring the two membranes together. This is illustrated schematically in Figure 1A. The N-helices (one from each subunit of the trimer) form highly conserved hydrophobic grooves into which the C-helices pack. It is generally agreed that formation of this six-helix structure is required for membrane-fusion to occur.

The importance of trimer-of-hairpins formation for HIV-1 entry led to the hypothesis that the C-terminal region of gp41 might serve as a target for potential membrane-fusion inhibitors. C-peptides have been shown to inhibit HIV-1 entry into cells, with IC_{50} values as low as 1 nM in vitro (C. T. Wild et al., Proc. Natl. Acad. Sci. USA 91, 9770-9774 (1994); D. C. Chan et al., Proc. Natl. Acad. Sci. USA 95, 15613-15617 (1998)). Evidence suggests that C-peptides work in a dominant-negative fashion by binding to the N-peptide region and disrupting trimer-of-hairpins formation. If the C-terminal region is accessible (at least transiently) prior to formation of the trimer-of-hairpins, then it is reasonable to expect that agents that bind to this region of gp41 N-terminal will prevent membrane fusion. Consistent with this notion, peptides derived from the gp41 N-terminal region (referred to as N-peptides) are modest inhibitors of HIV-1 membrane fusion. The inhibitory mechanism of N-peptides has not been determined, in part because these peptides have a strong tendency to aggregate.

Applicants reasoned that a single soluble molecule that contains a folded N-helical core and two of the three C-helices of the core trimer-of-hairpins would be highly stable and would bind a single C-peptide with high affinity. As described herein, the hypothesis that the C-peptide region of gp41 is a target for inhibition of

HIV-1 entry has been tested. Results of the assessment, also described herein, have shown that Five-Helix, which binds the C-peptide region of gp41, shows potent inhibitory activity against HIV-1 and against HIV-1 variants containing a diverse set of envelope proteins. These results point to the C-peptide region of HIV gp41 as a viable target to inhibit the formation of the trimer-of-hairpins, which is required for membrane fusion (and, thus, HIV infection of cells) to occur.

Described herein are results that show that a protein that binds to the C-peptide region of gp41 inhibits HIV entry into cells. Such proteins are inhibitors of HIV and serve as the basis for development of additional anti-HIV agents. They might also be used for generating a neutralizing antibody response that targets the N-terminal region of the gp41 ectodomain.

Five-Helix, as the proteins are designated, takes advantage of the binding properties of the N-helix peptide coiled coil while minimizing the tendency of the N-peptides to aggregate. In one embodiment of Five-Helix, five of the six helices that make up the core of the gp41 trimer-of-hairpins structure are connected with (joined by) short peptide linkers. (See Figure 1A.) In this embodiment, Five-Helix lacks a third C-peptide helix, thus creating a vacancy in order to create a high-affinity binding site for the C-terminal region of gp41. In further embodiments of Five-Helix, the three N-peptide helices and more than two (but less than three complete) C-peptide helices are connected with short peptide linkers. In these embodiments, the three N-peptide helices, two complete C-peptide helices and a portion of the third C-peptide helix are connected with peptide linkers. The portion of the third C-helix can be as few as one amino acid residue of the third C-helix or any number of additional amino acid residues of the helix up to, but not including, all of the amino acid residues of the helix. Five-Helix protein of the present invention is soluble under physiological conditions.

The core of the trimer-of-hairpins, as formed by individual N- and C-peptides, is already quite stable, with a melting temperature of 65 °C. Applicants have shown that if 5 of the 6 helices are covalently joined to form a 5-Helix protein, the stability of the core is further increased (the stability is greater than the stability of the 6-Helix core). Under physiological conditions, Five-Helix is folded, soluble,

and stable. It has an α -helical content in close agreement with the value predicted from the design. (See Figures 2A and 2B.) In affinity-interaction experiments, Five-Helix interacts strongly and specifically with epitope-tagged C-peptides. (See Figure 2C.) This interaction induces a helical conformation in the bound C-peptide as judged by the difference in circular dichroism before and after mixing. (See Figure 2D.) These properties are consistent with the intended design of Five-Helix.

Five-Helix potentially inhibits HIV-1 membrane fusion (nanomolar IC_{50}) as measured by viral infectivity and cell-cell fusion assays. (See Figures 3A and 3B.) In contrast, a control protein, denoted Six-Helix, in which the C-peptide binding site is occupied by an attached C-peptide (i.e., all six helices that constitute the gp41 trimer-of-hairpins have been linked into a single polypeptide, as described in Example 1), does not have appreciable inhibitory activity. (See Figure 3A and Figures 8 and 9). Similarly, a Five-Helix variant, denoted Five-Helix(D4), in which the C-peptide binding site is disrupted by mutation of four interface residues (V549, L556, Q563 and V570) to Asp, does not block the membrane fusion event even at 1 μ M. (See Example 3 and Figure 3A.) These results support the conclusion that C-peptide binding is the key determinant of antiviral activity in Five-Helix.

The inhibitory activities of 5-Helix and C-peptides are expected to be antagonistic: when 5-Helix binds C-peptide, the amino acid residues thought to be responsible for the antiviral activities of each inhibitor are buried in the binding interface. Indeed, mixtures of 5-Helix and C34 [a well characterized and potent peptide inhibitor with an IC_{50} of approximately 1 nM] display a dose-dependent antagonistic effect (Figure 3B). In the presence of 5-Helix, high-potency inhibition by C34 is only observed when the peptide is in stoichiometric excess (Figure 3B).

Five-Helix inhibits infection by viruses pseudotyped with a variety of HIV-1 envelope proteins (from clades A, B, and D) with similar potency (Figure 3D). This broad-spectrum inhibition likely reflects the highly conserved interface between the N- and C-terminal regions within the gp41 trimer-of-hairpins structure (Figure 4). The residues in the C-peptide region of gp41 that are expected to make contact with 5-Helix are highly conserved in HIV-1, HIV-2, and SIV (Figure 4).

As a potent, broad-spectrum inhibitor of viral entry, Five-Helix may serve as the basis for the development of a new class of therapeutic agents against HIV-1. Although they typically require parenteral administration, protein-based therapeutics can be practical, as exemplified by insulin, growth hormone, tissue plasminogen activator, granulocyte-colony stimulating factor, and erythropoietin. Alternatively, Five-Helix could be expressed endogenously (e.g., via gene therapy) with secretion into the bloodstream. If Five-Helix were expressed endogenously in HIV-infected cells, it could inhibit intracellular folding and transport of gp160. Five-Helix, Five-Helix (D4), and Six-Helix are also potential reagents for small-molecule drug-screening purposes. Five-Helix offers a great deal of flexibility in the design of variants with better therapeutic characteristics. In principle, Five-Helix can be modified extensively, except at its C-peptide binding site, to alter its immunogenic, antigenic, bioavailability, or inhibitory properties. For example, the C-peptide binding site might be lengthened, shortened, or shifted in the gp41 sequence in order to optimize inhibitory potency by targeting different regions of the gp41 ectodomain.

It would be desirable to generate neutralizing antibodies that mimic the binding properties of Five-Helix. The broadly neutralizing ability of Five-Helix most likely stems from its interaction with the highly conserved residues in the C-peptide region of gp41 (Figure 4). Unstructured C-peptide immunogens may not elicit broadly neutralizing antibodies because the linear sequence of the gp41 C-peptide region is variable among different HIV-1 strains. Such unstructured C-peptides do not have a long region of conserved amino acids residues. Rather, conserved amino acid residues and nonconserved residues are interspersed. However, constraining C-peptides or C-peptide analogues into a helical conformation (e.g., as in the C-peptide region when it binds Five-Helix) may lead to useful immunogens in the effort to develop an AIDS vaccine. Figure 4 is a helical wheel diagram depicting the interaction of Five-Helix with the C-peptide region of gp 41. As shown, on the helical wheel, the whole "face" is comprised of conserved or identical amino acid residues. As also shown, there is a high degree of conservation in the a and d positions of the C-peptide region of HIV gp41. Peptides from the C-terminal region of the gp41 ectodomain constrained in such a manner

-12-

that they present highly conserved amino acid residues on a single face of the molecule (such as in positions a, d and e in Figure 4) can be produced. They can be used as immunogens to produce antibodies that will presumably bind those amino acid residues in the corresponding unconstrained peptide (C-peptide region of HIV gp41) and, thus, mimic the binding characteristics of Five-Helix. For example, antibodies that bind some or all of the highly conserved (identical and/or conserved) amino acid residues in C38 (see Figure 4) can be produced. Such antibodies, which mimic the binding of Five-Helix, will work, in effect, as a preventive or vaccine by reducing or preventing the activity (binding) of Five-Helix. Such antibodies to constrained peptides from the C-terminal region of HIV gp41 ectodomain are a subject of this invention.

Intriguingly, the epitope for 2F5, the only known human monoclonal antibody directed against gp41 with broad neutralizing activity, is located immediately C-terminal to the C-peptide region targeted by Five-Helix (T. Muster, et al., J. Virol. 67, 6642-6647 (1993); M. Purtscher, et al., AIDS 10, 587-593 (1996)). The core of the 2F5 epitope (Leu-Asp-Lys-Trp; residues 663-666 in the HIV HXB2 gp 160 sequence) is highly conserved (81% identity) across the same set of HIV-1, HIV-2, and SIV isolates used to generate Figure 4. However, some HIV-1 escape variants to 2F5 neutralization do not contain mutations in the epitope sequence, suggesting that inhibition by 2F5 may involve recognition of additional determinants. The conformation of the 2F5-bound epitope remains unknown, but antibodies elicited with fragments of gp41 containing this sequence do not possess significant virus-neutralizing activity (T. Muster, et al., J. Virol. 68, 4031-4034 (1994); L. Eckhart, et al., J. Gen. Virol. 77, 2001-2008 (1996)). It remains to be seen if 2F5 inhibits infection by interfering with trimer-of-hairpins formation.

Further, Five-Helix itself is a vaccine candidate. The possibility of eliciting an antibody response against transiently exposed conformations of proteins involved in HIV-1 fusion has been suggested (R. A. LaCasse, et al., Science 283, 357-362 (1999)). One possible well-defined target is the N-terminal coiled coil that is exposed in the prehairpin intermediate (D. M. Eckert, et al., Cell 99, 103-115

-13-

(1999)). A 5-Helix-like intermediate may be exposed during the fusion process, and, in this case, antibodies directed against 5-Helix may inhibit viral entry.

Results described herein point to the C-peptide region of HIV-1 gp41 as a viable target to inhibit the formation of the trimer-of-hairpins. Structural and
5 computational methods predict similar trimer-of-hairpins motifs for viruses in many diverse families, including orthomyxoviridae, paramyxoviridae, filoviridae, retroviridae, and others. Moreover, in some of these cases, inhibition of viral entry by peptides analogous to the C-peptides of gp41 has been demonstrated. Thus, the Five-helix design approach may offer a widely applicable strategy for inhibiting viral
10 infections.

In addition, Five-Helix provides a means to study a formed C-peptide binding site in detail, which cannot be done with aggregable N-peptides. The exposed C-peptide binding site in this Five-Helix molecule is useful to identify or design molecules that bind to the N-helical core of gp41 and can be further assessed,
15 using known methods, for their ability to inhibit fusion of the HIV membrane with the membrane of a mammalian cell, such as a human cell, thus inhibiting (reducing or preventing) infection of the cell. Further, Five-Helix can be assessed for its ability to bind to the C-helical region of gp41 and inhibit its function. The N-helical core of gp41 is highly conserved (in terms of amino acid composition) and, thus, it is
20 likely that 5-Helix and variants thereof will be broadly neutralizing against a variety of clinical HIV strains and, thus, useful therapeutically.

The Five-Helix protein, which is based upon the known structure of the gp41 ectodomain, consists, in one embodiment, of three N-peptides and two C-peptides covalently linked and arranged to fold into a substantial part of the N-helical core
25 with two of the three C-helix binding sites occupied by C-peptide. The remaining C-peptide binding site of the N-peptide is exposed. The site exposes an epitope that is 40 amino acids in length. In addition, it is expected that the backbone atoms of the site are rigidly held in a structured conformation, as the N-peptide core is locked into place by the outer two C-peptides.

30 In single letter amino acid code, the amino acid sequence of one embodiment of Five-Helix is the following:

-14-

MQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILAGGSGGHTTWM
 EWDREINNYTSLIHSLEESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLR
 AIEAQQHLLQLTVWGIKQLQARILAGGSGGHTTWMEWDREINNYTSLIHSLE
 EESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGI
 5 KQLQARILAGGR (SEQ ID NO.: 1).

In single letter amino acid code, the amino acid sequence of 6-Helix is the following:

MQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILAGGSGGHTTWM
 EWDREINNYTSLIHSLEESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLR
 10 AIEAQQHLLQLTVWGIKQLQARILAGGSGGHTTWMEWDREINNYTSLIHSLE
 EESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGI
 KQLQARILAGGRGGHTTWMEWDREINNYTSLIHSLEESQNQQEKNEQELLG
 GHHHHHH (SEQ ID NO.: 2).

Five-Helix protein can be produced by a variety of methods. For example, it
 15 can be produced, as described in Example 1, from a larger protein, such as 6-Helix,
 by enzymatic (trypsin) digestion. Alternatively, it can be produced, using known
 methods and expression systems, by expressing Five-Helix protein-encoding DNA,
 which can be a single DNA that encodes the entire Five-Helix protein or two or
 more DNA "units", each of which encodes a portion (e.g., one or more N helices,
 20 one or more C helices) of N-Helix protein. The yield of expression and purification
 of Five-Helix can be significantly improved by direct expression of the Five-Helix
 gene in an appropriate host cell, such as *E. coli*. In this approach, the Five-Helix
 gene encodes the residues present in the final Five-Helix protein. A C-terminal His-
 tag can be attached to facilitate purification (with or without a protease cleavage site
 25 to later remove the tag). The protein can then be used directly without the
 proteolytic cleavage and unfolding steps required for producing Five-Helix starting
 from Six-Helix. This Five-Helix molecule may be expressed as a folded active
 molecule, allowing its use in biological selections or screens for optimizing its
 properties. Alternatively, protein synthetic methods can be used to produce Five-
 30 Helix protein. The five helices of Five-Helix can be joined covalently (such as by
 means of a linker of at least one (one or more) amino acid residues) or by other

-15-

means which results in formation of a protein which is stable under physiological conditions and is correctly folded such that the remaining surface of Five Helix is presented so that it is available to bind C34 peptide. In the embodiments in which there are three N-helices and more than two (but less than three complete) C-helices, the helices can be similarly joined.

Five-Helix can have a wide variety of sequences, both in the N- and C-helix regions and in the linker components, and can be comprised of L-amino acid residues, D-amino acid residues or a combination of both L- and D-amino acid residues. The amino acids residues can be modified. Five-Helix can include amino acid residues in addition to those of the helices and linkers (e.g., to stabilize the molecule). It is likely that the Five-Helix described here can be altered to enhance stability and activity. Minor changes in the design of the loops connecting the N- and C-helices (both in length and composition) and the exact borders of the N- and C-helices are likely to have significant effects on the stability, yield, and activity of Five-Helix.

As currently constructed, Five-Helix exposes a C-peptide binding site encompassing 40 amino acids along the N-helical core. A strategy for exposing shorter segments of the C-peptide binding site on 5-helix (or related molecules) involves attaching a short C-peptide sequence onto the longer exposed epitope. A molecule of this type might aid in the development of drugs targeted specifically to a shorter epitope along the N-helical core. For instance, a single pocket region (similar to that found in IQN17; D. M. Eckert, et al., Cell 99, 103-115 (1999)) could be exposed in Five-Helix by binding a C-peptide that lacks the residues that bind there (the first 10 or so residues of C34). These short C-peptide sequences could be attached to Five-Helix through a variety of means, including covalent crosslinking or merely extending the sequence of Five-Helix to cover part of the exposed epitope.

Five-Helix is useful in a variety of contexts. As described herein, Five-Helix is a potent inhibitor of viral membrane fusion, and, thus, acts on the virus before it enters the cell (unlike current practical therapy) which acts on HIV-infected cells. Five-Helix is soluble and has been shown to be stable under the conditions described herein. It should also be possible to generate 5-Helix variants with an increased

molecular weight (by oligomerization or tethering to a large protein) to reduce the rate of kidney clearance. In addition, Five-Helix dimers can be made by disulfide crosslinking, to produce a molecule filtered to a lesser extent than the Five-Helix “monomer”. Thus, it is reasonable to expect that dimers might have an enhanced
5 bioavailability when compared to that of the C-peptides.

Five-Helix prevents virus from entering cells, unlike standard therapy that targets viral proteins after viral entry, and thus, Five-Helix can be used prophylactically to prevent infection or reduce the extent to which infection occurs. One use for such a therapeutic is in the event of a needlestick injury, such as might
10 occur in a hospital or in settings in which needles contaminated with HIV are shared. For example, an individual who is stuck with a needle and is or might be infected with HIV can receive a sufficient quantity of Five-Helix (therapeutically effective quantity) in one or more dose(s) in order to prevent or reduce HIV entry into cells. Five-Helix can be administered, for example, by intravenous or intramuscular
15 injection.

In one embodiment of the present invention, Five-Helix is used to reduce HIV infection in an individual. In this embodiment, Five-Helix is administered, either as Five-Helix itself or via expression of Five-Helix-encoding DNA in appropriate host cells or vectors, to an individual in sufficient quantity to reduce
20 (totally or partially) HIV infection of the individual’s cells. That is, a dose of Five-Helix sufficient to reduce HIV infection (an effective dose) is administered in such a manner (e.g., by injection, topical administration, intravenous route) that it inhibits (totally or partially) HIV entry into cells. In one embodiment, a gene therapy approach is used to provide the effective dose, by introducing cells that express
25 Five-Helix protein into an individual. Five-Helix can be administered to an individual who is HIV infected to reduce further infection, or to an uninfected individual to prevent infection or reduce the extent to which infection occurs.

Pharmaceutical compositions which comprise Five-Helix in an appropriate carrier (e.g., a physiologically acceptable buffer) are a subject of this invention.
30 They are useful for preventive and therapeutic purposes and can be administered via a variety of routes (e.g., injection, topical administration, intravenous route).

-17-

Five-Helix appears to present a single, intact C-helix binding site and, thus, is useful for screening for drugs that inhibit membrane fusion. Five-Helix exposes a larger, more rigid target for potential drug screens than does IQN17. The molecules 6-Helix and 5-Helix(D4) are a useful negative control in these studies.

5 The Five-Helix exposed epitope can also be used as an antigen for producing antibodies, particularly neutralizing antibodies using known methods. The antibodies can be monoclonal or polyclonal.

10 The serum stability of Five-Helix can be tested, using known methods, to ascertain its therapeutic potential. If Five-Helix is degraded, the most likely point of attack/degradation is the glycine/serine linker regions. In this case, different linker regions can be generated and tested (see below). The inhibitory ability of these anti-Five-Helix sera and ascites can be tested using standard fusion assays.

15 The outside surface of Five-Helix can be varied, for example, to enhance bioavailability, decrease toxicity and avoid immune clearance. Since Five-Helix exhibits potent inhibitory activity, whereas the 6-Helix bundle does not, it is the exposed groove, including the pocket region, that is responsible for inhibition. The rest of the molecule simply provides a scaffold for displaying the exposed groove. Therefore, this scaffold can be modified without adversely affecting the inhibitory activity of Five-Helix. Modification of the scaffold may provide several advantages.

20 First, it would facilitate procedures in which multiple administrations of Five-Helix are required. For example, when Five-Helix is used as an anti-HIV therapeutic agent, multiple doses might be required. After extended administration, individuals might develop antibodies to Five-Helix that are likely to increase its clearance from the body. The availability of multiple versions of 5-Helix would help to circumvent

25 this problem by evading pre-existing antibodies. Second, it may be possible to design versions of Five-Helix, for example by introducing glycosylation sites on the external surface, in which the scaffold is less immunogenic. For vaccine studies, this modification would help to bias the immune response toward the exposed groove as opposed to the scaffold.

30 The observation that binding the gp41 C-helical region prevents HIV infection suggests a strategy for constructing an HIV vaccine. Analogous to

-18-

inhibition of HIV by C-peptides, Five-Helix likely inhibits gp41 by binding to a fusion intermediate of gp41 called the prehairpin intermediate. Whereas the C-peptide inhibitors function by binding to the N-peptide region of this intermediate, Five-Helix likely functions by binding to the C-peptide region. These considerations
5 strongly suggest that the C-peptide region of gp41 is a good drug target for the development of HIV entry inhibitors. Moreover, it may be possible to use C-peptide-based constructs as immunogens to elicit neutralizing antibodies. In the case of Five-Helix, the target of inhibition is a helical conformation of the C-peptide region, but reagents targeting other conformations of the C-peptide region may also
10 have inhibitory activity.

Recent vaccine studies (R.A. LaCasse, et al., Science 283, 357-362 (1999)) suggest that intermediates of the envelope-mediated fusion process can elicit strongly neutralizing antibodies. Antibodies to such fusion intermediates would target conserved regions of the envelope proteins and therefore would be likely to
15 neutralize a broad range of viral strains. Antibodies to the C-peptide region would target a region that is highly conserved and critical to the fusion process.

The trimer-of-hairpins is a common feature of many viral membrane fusion proteins. It has been observed in crystal structures of Influenza, Ebola, SV5 (simian parainfluenza virus 5), and RSV (human respiratory syncytial virus). In addition,
20 many other members of the retrovirus, paramyxovirus, and filovirus families are predicted to contain this motif. A similar structure has been observed in the associated vertebrate vesicle fusion proteins. The basic strategy described herein can be applied to any of these systems in order to inhibit fusion. One subject of this invention is a method of inhibiting formation of the trimer-of-hairpins of an
25 enveloped virus (a virus that comprises a viral envelope protein) by contacting the virus with a drug that binds a viral envelope protein (e.g., the C peptide region of a viral envelope protein) and inhibits formation of the trimer-of-hairpins of the enveloped protein.

The present invention is illustrated by the following examples, which are not
30 intended to be limiting in any way.

Example 1. Production of 5-Helix

- The design of 5-Helix was based on the N36/C34 six-helix bundle crystal structure (D. C. Chan, et al., Cell 89, 263-273 (1997)). For the 5-Helix protein, each peptide region was extended (compared with N36 and C34) by three residues on its N-terminus and one residue on its C-terminus, generating the final N40 and C38 segments (representing residues 543-582 and 625-662 of HIV-1 HXB2 gp160, respectively). Three N40 and two C38 segments were joined using a -GGSGG-linker after N40 and a -GSSGG-linker after C38. All constructs include an N-terminal Met for translation initiation. Two distinct 5-Helix proteins that differ only at their C-termini were generated for this study: (i) His-tagged 5-Helix, which ends in -GG(H)6, and (ii) untagged 5-Helix, which ends in -GGR. In addition, a third construct, denoted 6-Helix, was generated in which the 5-Helix backbone was connected to the His-tagged C-peptide, C37-H6 (see Example 4), through a trypsin-cleavable linker (-GGR-) (see Figures 10 and 11A-11C).
- All DNA constructs were assembled from PCR cassettes sequentially cloned into the pAED4 vector [D. S. Doering, P. Matsudaira, Biochemistry 35, 12677-12685 (1996)] using *E. coli* XL1-Blue (recA- strain, Stratagene). All proteins were recombinantly expressed in *E. coli* strain RP3098 grown in 2xYT to an OD (590 nm) between 0.5-0.7 before induction with IPTG (0.4 mM) for 3 hours. Bacterial pellets were resuspended in Tris/NaCl buffers (Qiaexpressionist booklet, March 1999, Qiagen) supplemented with Complete EDTA-free protease inhibitor tablets (Roche), and subsequently frozen at -20°C until the day of purification. Thawed resuspensions were lysed (sonication or French press) and centrifuged (35,000 x g for 30 minutes) to separate the soluble fraction from inclusion bodies.
- His-tagged 5-Helix (generated from plasmid p-5HelixH6) was purified directly from the inclusion bodies resuspended in 8 M urea in TBS (50 mM Tris, pH 8.0, 100 mM NaCl) and 10 mM imidazole. The mixture was clarified by centrifugation (35,000 x g for 30 minutes) before binding to a Ni-NTA agarose (Qiagen) column at room temperature. Protein was eluted in 6 M urea/TBS/100 mM imidazole in 40 ml (~5 column volumes). The protein was refolded by slow dripping into a one liter, stirred solution of 20 mM Tris (pH 8.0) at room

-20-

temperature. Refolded protein was then reconcentrated by passage over a Ni-NTA agarose column and eluted with 20 ml (~2 column volumes) of 100 mM imidazole in TBS.

Untagged 5-Helix was produced via proteolysis of 6-Helix (see below) to
5 generate a 5-Helix/C37-H6 complex. Following digestion with trypsin (1:200 weight ratio in TBS at room temperature for 1 hour, Sigma), the 5-Helix/C37-H6 complex was bound to Ni-NTA agarose and washed extensively to remove excess trypsin. The beads were resuspended in 8 M GuHCl/TBS and heated (70°C) in order to denature the complex. The nonbinding fraction, containing denatured 5-Helix,
10 was sequentially dialyzed into 8 M urea/20 mM Tris, pH 8.0 (4 hours at room temperature) and 4 M urea/20 mM Tris, pH 8.0 (overnight at 4°C). The protein was loaded onto a DEAE column (Fastflow, Pharmacia) and a reverse urea gradient (4 M to 0 M urea in 20 mM Tris, pH 8.0) was run over 20 column volumes in 4 hours at room temperature. The protein was eluted from the DEAE resin using a NaCl
15 gradient (0 to 300 mM) in 20 mM Tris, pH 8.0 (10 column volumes). 6-Helix (generated from plasmid p-6Helix) was purified directly from the soluble fraction of the bacterial lysate. The solution was passed over Ni-NTA agarose column and eluted with an imidazole gradient (10-250 mM) in TBS over 10 column volumes.

20 For all proteins, monomers were separated from aggregates by gel filtration (Sephacryl S200 HR or Superdex 75) in TBS. The proteins were >95% pure as judged by SDS-PAGE and can be concentrated to at least 3 mg/ml. The concentrations of all peptides and proteins were determined by absorbance at 280 nm in 6 M GuHCl [H. Edelhoch, Biochemistry 6, 1948-1954 (1967)].

25 **Example 2. Assessment of the specificity of 5-Helix/C-peptide interaction and of inhibition by 5-Helix of membrane fusion**

The specificity of 5-Helix/C-peptide interaction has been tested using a His-tagged C-peptide (C37-H6, independently expressed in *E. coli* and purified through reverse-phase HPLC) and Ni-agarose precipitation. In TBS with 30 µM of C37-H6 ,
30 16 µM of 5-Helix is completely precipitated by Ni-agarose. Addition of 150 µM

-21-

C34 (no His-tag, chemically synthesized and purified over HPLC) substantially reduces the amount of precipitated 5-Helix. The effective competition of C37-H6 and C34 indicates that 5-Helix binds C-peptide in a specific manner. The CD experiments and competitive binding assays suggest that 5-Helix folds into the predicted conformation. That is, the results support the prediction that 5-Helix contains an exposed C-peptide binding site.

Assays were carried out to assess the ability of 5-Helix to interact with the C-region of gp41 and inhibit function of the fusion protein. This inhibition of membrane fusion by 5-Helix and 6-Helix was assessed using a cell-based assay.

10 Proteins 5-Helix and 6-Helix are serially diluted in modified DMEM media with 5% FCS and aliquoted into slide chambers. HELA cells (4×10^4) expressing CD4 and coreceptors and containing a β -galactosidase gene under the control of the Tat promoter are added. CHO cells (2×10^4) expressing gp160 (precursor protein to gp120/gp41) and Tat are also added. The 400 μ l miniculture is incubated at 37 °C

15 for 8 to 24 hours; fused cells (syncytia) will transcribe and translate β -galactosidase. The cells are fixed in gluteraldehyde and exposed to X-gal/Fe solution for one hour. Syncytia that contain β -galactosidase turn blue-green. In this assay, 5-Helix demonstrates a potent inhibition of syncytia formation, with an IC_{50} of 10-20 nM; in one assay the IC_{50} was 13 nM. 6-Helix does not block fusion appreciably even at 1

20 μ M concentrations.

In order to verify the specificity of the 5-Helix exposed epitope as the inhibitory agent and to rule out a contaminant, mixing experiments with C-peptide have been performed. 5-Helix, at 200 nM concentration, is mixed with C34 at 100, 166, 190 and 210 nM. At the concentrations used, free 5-Helix and free C34 should

25 inhibit almost all of the syncytia in the miniculture. In 5-Helix/C34 mixes where C34 is in excess of the 5-Helix (i.e., at 210 nM) syncytia formation is not blocked in the 5-Helix/C34 mixes where C34 concentration is less than that of 5-Helix. By contrast, C34 in the presence of 6-Helix blocks all syncytia formation.

The inhibitory potentials of 5-Helix and 6-Helix have been reproduced in

30 viral fusion experiments. HIV, modified to contain a luciferase reporter gene, is mixed with human osteosarcoma (HOS) cells expressing CD4 and coreceptor in the

-22-

presence of diluted protein for 6 hours at 37 °C. The virus solution is replaced, and the HOS culture is incubated 48 hours more in fresh media. Luciferase activity is measured in a luminometer. In this assay, 5-Helix inhibits luciferase activity with an IC_{50} less than 10 nM. Again, 6-Helix shows no appreciable block up to 1 μ M (Figures 3A and 3C).

Example 3. Design and assesment of 5-Helix (D4)

In 5-Helix(D4), four highly conserved residues in the C-peptide binding site of His-tagged 5-Helix (Val549, Leu556, Gln563, and Val570) were mutated to Asp in the final (third) N40 segment. The construct [p-5Helix(D4)] was recombinantly expressed and purified in the same manner as the His-tagged 5-Helix. The His-tagged 5-Helix and 5-Helix(D4) proteins have the same ellipticity: for both, $[\theta]_{222} = -28,100 \pm 1500 \text{ deg cm}^2 \text{ dmol}^{-1}$ (~100% of the predicted helical content) at 4°C in TBS, and both proteins are extremely stable to thermal denaturation ($T_m > 98^\circ\text{C}$) in TBS, as well as to GuHCl chemical denaturation (C_m values ~6 M for 5-Helix(D4); ~7.2 M for the His-tagged 5-Helix) at 25°C. The slightly decreased stability of 5-Helix(D4) likely reflects the low helical propensity and charge of the Asp residues which, in this context, are placed within a predominantly hydrophobic groove on the surface of 5-Helix.

Example 4. His-tagged C-peptide C37-H6

Peptide C37-H6 is a His-tagged C-peptide of the following sequence:
GGHTTWMEWDREINNYTSLIHSLIEESQNQQEKNEQELLGHHHHHH
 (SEQ ID NO.: 5). The peptide is derived from HIV-1 HXB2 residues 625-661 (underlined) and contains the entire C34 sequence (W628 to L661). C37-H6 is produced from the tryptic digestion of a recombinantly expressed construct, p4-NC1.1, consisting of one N40 segment joined to C37-H6 through a -GGR- linker. Following expression, NC1.1 is purified from the soluble fraction of bacterial lysates in the same manner as 6-Helix. Trypsin digestion (same conditions as for untagged 5-Helix) generates C37-H6, which is then purified to homogeneity by reverse phase HPLC using a Vydac C-18 column and a linear gradient of acetonitrile in water

-23-

containing 0.1% trifluoroacetic acid. The identity of C37-H6 was confirmed by mass spectrometry (MALDI-TOF, PerSeptive). Like C34, C37-H6 is a potent inhibitor of HIV-1 membrane fusion, with an $IC_{50} \approx 1$ nM in the cell-cell fusion assay.

5 The data in Figures 2A-2D were generated using the untagged version of 5-Helix, but similar results were obtained with the His-tagged version [see Example 3]. The CD (Aviv 62 DS) experiments were performed in TBS buffer unless otherwise stated. In Fig. 2B, the protein concentration was 1 mM for the TBS sample and 0.54 mM for the GuHCl/TBS sample. In Fig. 2D, a quartz mixing cell
10 (Helma) with 1 ml chambers (4.375 mm/chamber pathlength) was utilized. The polypeptides were at a concentration of 5.9 mM (5-Helix) and 6 mM (C37-H6) in 20 mM Tris, pH 8.0/250 mM NaCl before mixing.

The 5-Helix precipitation experiment (Fig. 2C) was performed in 20 ml TBS with 16 mM untagged 5-Helix, 30 mM His-tagged C37-H6, and/or 150 mM C34.

15 The solution was added to 10 ml of Ni-NTA agarose and incubated at room temperature for 10 minutes. After the unbound supernatant was removed, the beads were washed twice with 1 ml TBS and then eluted with 500 mM imidazole. The Ni-bound and unbound samples were run on a 16.5% Tris-Tricine polyacrylamide gel (Biorad) and stained with Gel-code Blue (Pierce).

20 Example 5. His-tagged 5-Helix

All data in Figures 3A-3C were generated using His-tagged 5-Helix (see Example 1). The cell-cell fusion assays (Figure 3B) were performed as described (D. C. Chan et al., Proc. Natl. Acad. Sci. USA 95, 15613-15617 (1998)). Inhibition of viral infectivity was studied using a recombinant luciferase reporter assay slightly
25 modified from that previously detailed (D.C. Chan, et al., Proc. Natl. Acad. Sci., USA, 95, 15613-15617 (1998)). Briefly, pseudotyped viruses were generated from 293T cells cotransfected with an envelope-deficient HIV-1 genome NL43LucR⁻E⁻ [B. K. Chen, et al., J. Virol. 68, 654-660 (1994)] and one of four gp160 expression vectors: pCMV-HXB2 (D.C. Chan, et al., Proc. Natl. Acad. Sci., USA, 95, 15613-
30 15617 (1998)), pEBB-JRFL (kindly provided by B. K. Chen), pSVIII-UG024.2, and

-24-

pSVIII-RW020.5. The plasmids pSVIII-UG024.2 and pSVIII-RW020.5 were obtained from the NIH AIDS Reagent Program (F. Gao, B. Hahn, and the DAIDS, NIAID) and code for envelope protein from primary HIV-1 isolates. Supernatants containing virus were prepared as previously described (D. C. Chan et al., Proc. Natl. Acad. Sci. USA 95, 15613-15617 (1998)) and used to infect either HOS-CD4 cells (HXB2 and UG024.2) or HOS-CD4-CCR5 cells (JRFL and RW020.5). Cells were obtained from the NIH AIDS Reagent Program (N. Landau). In Figure 3A, viral infectivity assays were performed in the standard 24-well format (D. C. Chan et al., Proc. Natl. Acad. Sci. USA 95, 15613-15617 (1998)). The data in Figure 3C were obtained from assays conducted in 96-well format: virus-containing supernatant (10 ml) and media (90 ml) were overlaid onto HOS cells at 50% confluency. Following two days of incubation at 37°C, the cells were harvested in 100 ml lysis buffer (Luciferase Assay System, Promega), of which 10 ml was analyzed per manufacturer's protocol. The IC_{50} values were calculated by fitting the 5-Helix titration data to a Langmuir function [normalized luciferase activity = $1/(1 + [5\text{-Helix}]/IC_{50})$].

While this invention has been particularly shown and described with references to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the scope of the invention encompassed by the appended claims.

-25-

CLAIMS

What is claimed is:

1. Five-Helix protein which is soluble under physiological conditions and comprises the three-N-helices and at least two, but not three complete, C-helices of the trimer of hairpin structure of HIV gp41, wherein the component helices are separated by linkers.
2. Five-Helix of claim 1 wherein the linker comprises at least one amino acid residue linker.
3. Five-Helix protein of claim 2 wherein the protein binds the C-peptide region of HIV gp41.
4. Five-Helix protein of claim 2 which includes the three N-helices and two C-helices of the trimer of hairpin structure of HIV gp41.
5. Five-Helix comprising SEQ ID No.: 1.
6. Six-Helix comprising the three N-helices and the three C-helices of the trimer hairpin structure of HIV gp41, wherein the component helices are separated by linkers.
7. Six-Helix protein comprising SEQ ID NO.: 2.
8. A method of identifying a compound or molecule that binds Five-Helix and inhibits HIV infection of mammalian cells, wherein the compound or molecule to be assessed is referred to as a candidate inhibitor, comprising combining a candidate inhibitor and Five-Helix, under conditions appropriate for binding of an inhibitor and Five-Helix to occur and determining if

-26-

binding occurs, wherein if binding occurs, the candidate inhibitor is a compound or molecule that binds Five-Helix.

9. The method of claim 8 further comprising determining if the compound or molecule that binds Five-Helix inhibits HIV infection of mammalian cells in a cell-based assay.
10. A method of eliciting an immune response to HIV in an individual, comprising introducing, by an appropriate route, a composition comprising 5-Helix and a physiologically acceptable carrier, in a dose sufficient to elicit the immune response in the individual.
11. Five-Helix that binds the C-peptide region of HIV gp41 and inhibits HIV infection of mammalian cells.
12. Five-Helix of claim 9 that inhibits HIV infection of human cells.
13. Five-Helix that interferes with formation of the HIV gp41 trimer of hairpin structure and inhibits HIV infection of cells.
14. Five-Helix that inhibits fusion of HIV and mammalian cell membranes, as measured by viral infectivity assay, cell-cell fusion assay or both.
15. Five-Helix of claim 14, wherein the mammalian cell membranes are human cell membranes.
16. Five-Helix of claim 14 wherein the protein inhibits HIV membrane fusion at nanomolar IC_{50} , as measured by viral infectivity assay or cell-cell fusion assay.

-27-

17. A method of eliciting a neutralizing anti-HIV response in an individual, comprising administering to the individual Five-Helix that binds the C-peptide region of HIV gp41.
18. A method of inhibiting HIV infection of cells in an individual, comprising
5 administering to the individual Five-Helix in sufficient quantity and by an appropriate route for Five-Helix to bind the C-terminal region of HIV gp41, whereby HIV membrane fusion and HIV infection of cells are inhibited.
19. A method of inhibiting fusion of HIV and human cell membranes in an individual, comprising administering to the individual a drug that inhibits
10 formation of the trimer-of-hairpins of HIV gp41, thereby inhibiting fusion of HIV and human cell membranes.
20. The method of claim 19 wherein the drug is Five-Helix or a neutralizing antibody that mimics the binding properties of Five-Helix.
21. A method of inhibiting formation of the trimer-of-hairpins of an enveloped
15 virus, comprising contacting the virus with a drug that binds a viral envelope protein and inhibits formation of the trimer-of-hairpins of the enveloped virus.
22. The method of claim 21 wherein the drug is Five-Helix or an antibody that mimics the binding properties of Five-Helix.
- 20 23. Five-Helix complex, wherein the complex comprises Five-Helix linked to a molecule that binds HIV envelope protein.
24. Five-Helix complex of claim 23, wherein the molecule that binds HIV envelope protein binds HIV gp120.

-28-

25. Five-Helix complex of claim 24 wherein the molecule that binds HIV gp120 is sCD4 or an antibody.
26. A method of inhibiting fusion of HIV and human cell membranes in an individual, comprising administering to the individual Five-Helix complex in
5 sufficient quantity and by an appropriate route for Five-Helix complex to bind HIV envelope protein, wherein Five-Helix complex comprises Five-Helix linked to a molecule that binds HIV envelope protein.
27. The method of claim 26, wherein the envelope protein is HIV gp120.
28. An isolated protein selected from the group consisting of :
- 10 (a) (SEQ ID NO.: 1)
(b) (SEQ ID NO.: 2)
(c) (SEQ ID NO.: 3)
(d) (SEQ ID NO.: 4)
(e) (SEQ ID NO.: 5)
15 (f) (SEQ ID NO.: 6)
(g) (SEQ ID NO.: 7)
(h) (SEQ ID NO.: 8)
(i) (SEQ ID NO.: 9)
(j) (SEQ ID NO.: 10)
20 (k) (SEQ ID NO.: 11)
(l) (SEQ ID NO.: 12).







Figure 1A

Figure 1B



C

	Ni-Bound	Unbound
5-Helix	+	+
C37-H6	+	+
C34	-	-

Figure 2C


A

Figure 3A


B


Figure 3B

C

Isolate	Clade	Coreceptor	IC ₅₀ (nM)
HXB2	B	CXCR4	1.9 ± 0.7
UG024.2	D	CXCR4	1.3 ± 0.2
JRFL	B	CCR5	5.6 ± 0.7
RW020.5	A	CCR5	5.9 ± 2.7

Figure 3C

C-peptide
Region of
gp41



5-Helix




Figure 4





Figure 5

6-Helix



5-Helix






Figure 6




a) view from top of molecule




b) side view

Figure 7

Figure 8





Sequence overview for 5-Helix paper:**(N40)** = QLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILA (SEQ ID NO:3)**(C38)** = HTTWMEWDREINNYTSLIHS�IEESQNQQEKNEQELLE (SEQ ID NO:4)**(C37-H6)** = GGHTTWMEWDREINNYTSLIHS�IEESQNQQEKNEQELLGGHHHHHH
(SEQ ID NO: 5)**(N40**)** = QLLSGIDQQQNNLDRAIEAQDHLLQLTDWGIKQLQARILA (SEQ ID NO:6)**6-Helix****M-(N40)-GGSGG-(C38)-GSSGG-(N40)-GGSGG-(C38)-GSSGG-(N40)-GGR-(C37-H6)**

MQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEW
 DREINNYTSLIHS�IEESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEA
 QQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEWDREINNYTSLIHS�IEESQNQ
 QEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARIL
 AGGRGGHTTWMEWDREINNYTSLIHS�IEESQNQQEKNEQELLGGHHHHHH
 (SEQ ID NO:2)

5-Helix-R
(Untagged)
M-(N40)-GGSGG-(C38)-GSSGG-(N40)-GGSGG-(C38)-GSSGG-(N40)-GGR

MQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEW
 DREINNYTSLIHS�IEESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEA
 QQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEWDREINNYTSLIHS�IEESQNQ
 QEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARIL
 AGGR
 (SEQ ID NO:1)

Figure 11A

His-tagged 5-Helix

M-(N40)-GGSGG-(C38)-GSSGG-(N40)-GGSGG-(C38)-GSSGG-(N40)-GG(H)₆

MQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEW
DREINNYTSLIHSLIEESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEA
QQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEWDREINNYTSLIHSLIEESQNQ
QEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARIL
AGGHHHHHH (SEQ ID NO: 7)

5-Helix (D4)

M-(N40)-GGSGG-(C38)-GSSGG-(N40)-GGSGG-(C38)-GSSGG-(N40)-GG(H)₆**

MQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEW
DREINNYTSLIHSLIEESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEA
QQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEWDREINNYTSLIHSLIEESQNQ
QEKNEQELLEGGSSGGQLLSGIDQQQNNLDRAIEAQDHLLQLTDWGIKQLQARIL
AGGHHHHHH (SEQ ID NO: 8)

5-Helix-H6-GC

M-(N40)-GGSGG-(C38)-GSSGG-(N40)-GGSGG-(C38)-GSSGG-(N40)-GG(H)₆-GC

MQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEW
DREINNYTSLIHSLIEESQNQQEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEA
QQHLLQLTVWGIKQLQARILAGGSGGGHTTWMEWDREINNYTSLIHSLIEESQNQ
QEKNEQELLEGGSSGGQLLSGIVQQQNNLLRAIEAQQHLLQLTVWGIKQLQARIL
AGGHHHHHHGC (SEQ ID NO: 9)

Figure 11B

C-peptides

C34

WMEWDREINNYTSLIHSLEESQNQQEKNEQELL (SEQ ID NO:10)

C37-H6

GGHTTWMEWDREINNYTSLIHSLEESQNQQEKNEQELLGGHHHHHH
(SEQ ID NO:5)

Figure 11C